

**Computational Fluid Dynamics (AE/ME 339)**  
**MAEEM Dept., UMR, Fall 2001**

Home Work Problem 8

This problem involves free convection flow at a heated vertical wall and the numerical solution of the simplified non-dimensionalized equations. Use the explicit method as discussed in class.

Solve for the dependent variables  $U_{\text{bar}}$ ,  $V_{\text{bar}}$ , and  $\theta$ . Use the initial and boundary conditions are follows:

$\tau = 0$ :

$$U_{\text{bar}} = V_{\text{bar}} = 0, \theta = 0.$$

$\tau > 0$ :

$$\xi = 0: \quad U_{\text{bar}} = V_{\text{bar}} = 0, \theta = 0.$$

$$\eta = 0: \quad U_{\text{bar}} = V_{\text{bar}} = 0, \theta = 1.$$

$$\eta = \text{infinity}: \quad U_{\text{bar}} = V_{\text{bar}} = 0, \theta = 0.$$

Use the following for dimensional calculations:  $T_w = 50 \text{ C}$ , and  $T_{\text{infty}} = 20 \text{ C}$ .

Assume that the coefficient of thermal expansion is constant and is given by the following expression.

$$\beta = \frac{1}{\left(\frac{T_w + T_{\infty}}{2} + 273\right)} \dots (1/K)$$

The time step must satisfy the approximate stability criterion given below.

$$\frac{2\Delta\tau}{(\Delta\eta)^2} \leq 1$$

For the finite domain for your calculation choose  $\xi_{\text{max}} = 100$ , and  $\eta_{\text{max}} = 25$ .


Use  $\Delta\xi = 10$ ,  $\Delta\eta = 2.5$  and  $\Delta\tau = 0.5$ .

Calculate the variables at  $\tau = 80$ . Show the result in tabular form as indicated below.

Plot the distributions of the dimensional velocities and temperature along a horizontal line where ( $\xi = \xi_{\text{max}}/2.0$ ).

Write a short note describing the procedure, details of the calculations and discussion of the results.

Attach a listing of your computer program.



Make 3 tables for U, V, and theta values for the nodes.  
Choose print interval such that you show only 10 values  
in each directions at equally spaced nodes.