

WHITNEY MAPS—A NON-METRIC CASE

BY

JANUSZ J. CHARATONIK (WROCLAW AND MÉXICO) AND
 WŁODZIMIERZ J. CHARATONIK (WROCLAW AND ROLLA, MO)

Abstract. It is shown that there is no Whitney map on the hyperspace 2^X for non-metrizable Hausdorff compact spaces X . Examples are presented of non-metrizable continua X which admit and ones which do not admit a Whitney map for $C(X)$.

Given a Hausdorff compact space X , we consider the space 2^X of all non-empty compact subsets of X equipped with the Vietoris topology. Any subspace $H(X)$ of the space 2^X is called a *hyperspace* of X . In particular $F_n(X)$ stands for the family of all non-empty subsets of X of cardinality at most n (where $n \in \mathbb{N}$), and $C(X)$ denotes the hyperspace of subcontinua of X (i.e., of connected members of 2^X). The reader is referred to [4] and [5] for needed information on hyperspaces.

A continuum X containing two points a and b is called an *arc* (from a to b) provided that each point of $X \setminus \{a, b\}$ separates a and b in X . We write ab to denote an arc with end points a and b . Note that an arc is metrizable if and only if it is homeomorphic to the closed unit interval $[0, 1]$.

Given a Hausdorff compact space X and its hyperspace $H(X)$, by a *Whitney map* for $H(X)$ we mean a mapping $\mu : H(X) \rightarrow ab$ such that

$$(0.1) \mu(\{x\}) = a \text{ for each point } x \in X;$$

$$(0.2) A \subsetneq B \text{ implies } \mu(A) < \mu(B).$$

When X is a compact metric space, then a Whitney map for 2^X or $C(X)$ does always exist, and several constructions of such mappings are known: see e.g. [5, 0.50.1–0.50.3, pp. 25–26] or [4, Theorem 13.4, p. 107; Exercises 13.5–13.8, pp. 108–109].

The following theorem answers a question of Robert Heath asked during a private conversation with the second named author.

THEOREM 1. *The following conditions are equivalent for a Hausdorff compact space X :*

2000 *Mathematics Subject Classification*: 54B20, 54C99, 54E35, 54F15.

Key words and phrases: continuum, hyperspace, metrizable, Whitney map.

- (1.1) X is metrizable;
 (1.2) there exists a Whitney map from the hyperspace 2^X onto an arc;
 (1.3) there exists a Whitney map from the hyperspace $F_2(X)$ onto an arc.

Proof. As stated above, the implication from (1.1) to (1.2) is well known. The one from (1.2) to (1.3) is obvious. It remains to show that (1.3) implies (1.1). So, assume (1.3). Let $\mu : F_2(X) \rightarrow ab$ be a Whitney map. Let $\{x_1, x_2, \dots\}$ be a countable infinite subset of X . Then there is a cluster point x_0 of this set. Continuity of the mapping $x \mapsto \mu(\{x_0, x\})$ implies that the family $B_n = \{[a\mu(\{x_0, x_n\})] : x_n \neq x_0\}$, $n \in \mathbb{N}$, of half-open intervals is a countable local basis for ab at a . Define $f : X \times X \rightarrow ab$ by $f(\langle x, y \rangle) = \mu(\{x, y\})$. Then the diagonal Δ of $X \times X$ is a G_δ -subset since $\Delta = \bigcap \{f^{-1}(B_n) : n \in \mathbb{N}\}$. Therefore X is metrizable by [1, Exercise 4.2.B, p. 264]. ■

COROLLARY 2. *If a Hausdorff compact space is non-metrizable, then there is no Whitney map for the hyperspace 2^X .*

For any dendron (i.e., a Hausdorff continuum such that any two of its distinct points are separated by a third one) X there is a canonical embedding of $F_2(X)$ in $C(X)$ (which maps any pair $\{x, y\}$ with $x \neq y$ to the unique arc xy). Therefore we get the following.

OBSERVATION 3. *For any non-metrizable dendron X there is no Whitney map from the hyperspace $C(X)$. In particular, there is no Whitney map from the hyperspace $C(X)$ for any non-metrizable arc X .*

The assumption that X is a dendron is essential in the above observation. To see this recall the following example K (see [2] and [3]).

Let D be a totally disconnected Hausdorff compact space without isolated points (in the metric case only the Cantor set has these properties), and let $f : D \rightarrow D$ be a homeomorphism such that for each $x \in D$ the orbits of x , i.e., the sets $\{f^n(x) : n \in \{0\} \cup \mathbb{N}\}$ and $\{f^{-n}(x) : n \in \{0\} \cup \mathbb{N}\}$ are dense in D . In the product $D \times [0, 1]$ identify $\langle x, 0 \rangle$ with $\langle f(x), 1 \rangle$ for each $x \in D$. Let K be the quotient space. It is shown in [3] that each proper subcontinuum of K is a metric arc and that covering dimension of K is one.

Observe that using the natural flow on K one can easily define, for any arc A in K , the length $\ell(A)$ of A . Then a Whitney map $\mu : C(K) \rightarrow [0, 1]$ can be defined as follows:

$$\mu(A) = \frac{\ell(A)}{1 + \ell(A)} \quad \text{for } A \in C(K) \setminus \{K\}, \text{ and } \mu(K) = 1.$$

PROBLEM 4. Characterize non-metrizable continua X for which there exists a Whitney map for $C(X)$.

Acknowledgements. We would like to express our appreciation to the referee, who has not only read the manuscript very carefully, but also solved two main problems and simplified some of the proofs of the first version of the paper. The referee certainly deserves to be the coauthor of the paper, but decided to remain anonymous. Thanking the referee for his job, we underline that we do not consider a note at the end of the paper to be a sufficient appreciation of the referee's contribution.

REFERENCES

- [1] R. Engelking, *General Topology*, Heldermann, Berlin, 1989.
- [2] A. Gutek, *A generalization of solenoids*, in: *Topology (Budapest, 1978)*, Colloq. Math. Soc. János Bolyai 23, North-Holland, Amsterdam, 1980, 547–554.
- [3] A. Gutek and C. L. Hagopian, *A nonmetric indecomposable homogeneous continuum every proper subcontinuum of which is an arc*, *Proc. Amer. Math. Soc.* 86 (1982), 169–172.
- [4] A. Illanes and S. B. Nadler, Jr., *Hyperspaces*, Dekker, New York, 1999.
- [5] S. B. Nadler, Jr., *Hyperspaces of Sets*, Dekker, New York, 1978.

Mathematical Institute
University of Wrocław
Pl. Grunwaldzki 2/4
50-384 Wrocław, Poland
E-mail: jjc@hera.math.uni.wroc.pl
wjcharat@hera.math.uni.wroc.pl

Department of Mathematics and Statistics
University of Missouri-Rolla
Rolla, MO 65409-0020, U.S.A.
E-mail: wjcharat@umr.edu

Instituto de Matemáticas
UNAM
Circuito Exterior
Ciudad Universitaria
04510 México, D.F., México
E-mail: jjc@gauss.matem.unam.mx

Received 3 July 1999;
revised 13 January 2000

(3799)