

68. Consider the series  $\sum_{n=0}^{\infty} (-1)^n x^{2n}$ .
- For which  $x \in \mathbb{R}$  does the series converge pointwise?
  - Does the series converge uniformly on  $[-1, 1]$ ?
  - Does the series converge in the  $L^2$  sense on  $[-1, 1]$ ?
69. Let  $\phi(x) = -1 - x$  if  $x \in [-1, 0)$  and  $\phi(x) = 1 - x$  if  $x \in (0, 1]$  and  $\phi(0) = 0$ .
- Find the full Fourier series of  $\phi$  in the interval  $(-1, 1)$ .
  - Graph the first five partial sums of the Fourier series (use a computer if you like).
  - Does the Fourier series converge in the mean square sense?
  - Does the Fourier series converge pointwise?
  - Does the Fourier series converge uniformly?
70. Let  $f$  be  $2\pi$ -periodic on  $\mathbb{R}$  and assume  $\int_{-\pi}^{\pi} |f(x)|^2 dx < \infty$ . Show that both  $\int_{-\pi}^{\pi} f(x) \cos(nx) dx$  and  $\int_{-\pi}^{\pi} f(x) \sin(nx) dx$  converge to zero as  $n$  tends to infinity.
71. Find  $\sum_{k=1}^n \sin(k\theta)$ .
72. For  $|a| < 1$ , find
- $\sum_{n=0}^{\infty} a^n \cos(n\theta)$ ;
  - $\sum_{n=1}^{\infty} a^n \sin(n\theta)$ .
73. Let  $e_n(x) = \frac{1}{\sqrt{2\pi}} e^{inx}$ , where  $x \in (-\pi, \pi)$ . Let  $f$  be continuous and  $2\pi$ -periodic on  $\mathbb{R}$ . Define  $f_m = \sum_{n=-m}^m (f, e_n) e_n$  and  $F_m = \frac{1}{m+1} \sum_{k=0}^m f_k$ .
- Establish the formula  $F_m(y) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) K_m(y-x) dx$ , where  $K_m(\theta) = \frac{1}{m+1} \sum_{k=0}^m \sum_{n=-k}^k e^{in\theta}$  is the so-called Fejér kernel.
  - Show that  $K_m(\theta) = \frac{1}{m+1} \frac{\sin^2 \frac{(m+1)\theta}{2}}{\sin^2 \frac{\theta}{2}}$  if  $\theta \neq 2\pi n$  for some  $n \in \mathbb{Z}$ .
  - Establish the formula  $F_m(y) - f(y) = \frac{1}{2\pi} \int_{y-\pi}^{y+\pi} [f(x) - f(y)] K_m(y-x) dx$ .
  - Draw the graph of  $K_m$  (use a computer if you like) for  $m \in \{2, 5, 8\}$ .
74. Consider the problem  $u_t = k u_{xx}$ ,  $0 < x < l$ ,  $u(x, 0) = \phi(x)$  with  $u_x(0, t) = u_x(l, t) = \frac{u(l, t) - u(0, t)}{l}$ .
- Assume that there are no negative eigenvalues and solve the problem.
  - Assume that limits can be taken term by term and find  $A, B$  with  $\lim_{t \rightarrow \infty} u(x, t) = A + Bx$ .
75. Fill in the details for the verification of the Gibbs phenomenon for  $f(x) = -1/2$  if  $-\pi < x < 0$ ,  $f(x) = 1/2$  if  $0 < x < \pi$  (pp 137/138).
76. Work on any two problems of your choice from Section 5.6.