

MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Chapter 12**

**Wiener Processes  
and Itô's Lemma**

11/13/2024 Math 5737/Econ 5337, Fall 2024 179

179

MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Definition 12.1**

A **stochastic process** is a family of random variables  $X=X(t)$ , where  $t$  could be integers or real numbers.

11/13/2024 Math 5737/Econ 5337, Fall 2024 180

180


MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Remark 12.2 (Normal RV)**

$X \sim N(0, 1)$   
 $P(X \leq 0) = 0.5$   
 $P(X \leq 0.2) = 0.5793$   
 $P(X \leq -0.2) = 0.4207$   
 $P(X \leq x) = 0.95$

$X \sim N(2, 4)$   
 $P(X \leq 2) = 0.5$   
 $P(X \leq 1) = 0.0540$   
 $P(X \leq x) = 0.95$

If  $X \sim N(\mu, \sigma)$ , then  $(X-\mu)/\sigma \sim N(0, 1)$



11/13/2024 Math 5737/Econ 5337, Fall 2024 181

181

MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Definition 12.3**

A stochastic process follows a **Wiener process** if

- the change  $\Delta W$  during a small period of time  $\Delta t$  is  $\Delta W = \epsilon(\Delta t)^{1/2}$ , where  $\epsilon \sim N(0, 1)$ ,
- the values of  $\Delta W$  for any two different short intervals of time  $\Delta t$  are **independent**.

11/13/2024 Math 5737/Econ 5337, Fall 2024 182

182

MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Example 12.4**

Suppose  $W$  follows a Wiener process and time is measured in years. Suppose the value of  $W$  is initially 25. What is the value of  $W$  at the end of one year? What is the value of  $W$  at the end of five years? Find  $P(W(1) > 26)$  and  $P(W(5) > 26)$ .

11/13/2024 Math 5737/Econ 5337, Fall 2024 183

183

MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Definition 12.5**

A stochastic process  $X$  follows a **generalized Wiener process** with drift rate  $a$  and variance rate  $b^2$  if

$$dX = adt + bdW,$$

where  $W$  is a Wiener process.

11/13/2024 Math 5737/Econ 5337, Fall 2024 184

184

MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

### Example 12.6

Assume that the cash position of a company, measured in thousands of dollars, follows a generalized Wiener process with a drift of 20 per year and a variance rate of 900 per year. Initially, the cash position is 50. Find the probabilities of negative cash positions after 1 year and after 3 months.

11/23/2024 Math 5737/Econ 5337, Fall 2024 185

185

MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

### Definition 12.7

An **Itô process** is a generalized Wiener process in which the parameters **a** and **b** are functions of **X** and **t**.

11/23/2024 Math 5737/Econ 5337, Fall 2024 186

186

MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

### Remark 12.8

Now we discuss the stochastic process usually assumed for the price  $S$  of a non-dividend-paying stock. A generalized Wiener process  $dS=adt+bdW$  is not appropriate as

- expected percentage change of  $S$  should remain constant, not its expected absolute change,
- uncertainty as to the size of future stock price movements should be proportional to the level of the stock price.

11/23/2024 Math 5737/Econ 5337, Fall 2024 187

187

MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

### Remark 12.8 (continued)

So we are using the Itô process

$$dS = \mu S dt + \sigma S dW,$$

where  $\mu$  is the **expected return** and  $\sigma$  is the **volatility** of the stock price. This model can be regarded as the limiting case of the random walk represented by binomial trees as the time step becomes smaller. The model is also known as **geometric Brownian motion**.

11/23/2024 Math 5737/Econ 5337, Fall 2024 188

188

MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

### Example 12.9

Consider a non-dividend-paying stock with volatility 30% (pa) providing expected return of 15%. Suppose the stock price is initially 100. Assuming the stock price follows GBM, what is the probability that the stock price after one week is more than 100?

11/23/2024 Math 5737/Econ 5337, Fall 2024 189

189

MISSOURI S&T MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

### Example 12.10

Here we discuss **Monte-Carlo simulation**. Let  $\mu=0.14$ ,  $\sigma=0.2$ ,  $\Delta t=0.01$ .

t	S(t)	$\epsilon$	$\Delta S(t)$
0	20.000	0.52	
0.01		1.44	
0.02		-0.86	
0.03		1.46	
0.04		-0.69	
0.05		-0.74	
0.06		0.21	
0.07		-1.1	
0.08		0.73	
0.09		1.16	
0.1		2.56	

11/23/2024 Math 5737/Econ 5337, Fall 2024 190

190

MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Theorem 12.11**

(Itô's lemma)

If  $dW$  is a WP,

$$dX = a(X,t)dt + b(X,t)dW,$$

and  $G = G(x,t)$ , then

$$dG = (G_x a + G_t + G_{xx} b^2 / 2)dt + G_x b dW.$$

11/13/2024 Math 5737/Econ 5337, Fall 2024 191

191

MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Corollary 12.12**

If  $dS = \mu S dt + \sigma S dW$  and  $G = G(s,t)$ , then

$$dG = (G_s \mu S + G_t + G_{ss} \sigma^2 S^2 / 2)dt + G_s \sigma S dW.$$

11/13/2024 Math 5737/Econ 5337, Fall 2024 192

192

MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Corollary 12.13**

If  $dS = \mu S dt + \sigma S dW$  and  $F$  is the forward price of a forward contract on the non-dividend-paying stock, then

$$dF = (\mu - r)F dt + \sigma F dW.$$

11/13/2024 Math 5737/Econ 5337, Fall 2024 193

193

MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Remark 12.14**

Note that  $F$  in Corollary 12.13 follows again geometric Brownian motion with the same variance rate as the stock price and a growth rate equal to the excess return of the stock price over the risk-free rate.

11/13/2024 Math 5737/Econ 5337, Fall 2024 194

194

MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY

**Corollary 12.15**

If  $dS = \mu S dt + \sigma S dW$  and  $G = \ln(S)$ , then

$$dG = (\mu - \sigma^2 / 2)dt + \sigma dW.$$

11/13/2024 Math 5737/Econ 5337, Fall 2024 195

195