

Lecture 27: Wave interference

- Superposition of waves
- Standing waves on a string
- Interference

Standing Waves

Two waves traveling in opposite directions with:

- same amplitude
- same wavelength λ (and thus same k)
- same frequency f (and thus same ω)

\Rightarrow same speed $v = \lambda f$.

$y_1(x, t) = A_0 \sin(kx - \omega t)$ traveling in the positive x-dir.

$y_2(x, t) = A_0 \sin(kx + \omega t)$ traveling in the negative x-dir.

Standing Waves

$$y_1(x, t) = A_0 \sin(kx - \omega t) \quad \text{traveling in the positive x-dir.}$$

$$y_2(x, t) = A_0 \sin(kx + \omega t) \quad \text{traveling in the negative x-dir.}$$

$$\sin a + \sin b = 2 \sin \frac{a+b}{2} \cos \frac{a-b}{2}$$

$$y_1 + y_2 = A_0 \cdot 2 \sin \frac{(kx - \omega t) + (kx + \omega t)}{2} \cos \frac{kx - \omega t - (kx + \omega t)}{2}$$

$$= 2A_0 \sin \frac{2kx}{2} \cos \frac{-2\omega t}{2}$$

$$y(x, t) = y_1 + y_2 = 2A_0 \sin(kx) \cos(\omega t)$$

Standing wave

Standing wave on a string


$$y(x, t) = A \sin(kx) \cos(\omega t)$$

String of length L
with fixed ends:

$$y(x = 0, t) = 0$$

$$y(x = L, t) = 0$$

Boundary Condition



$$\Rightarrow \sin(kL) = 0$$

$$kL = n\pi$$

With $k = \frac{2\pi}{\lambda}$:

$$\frac{2\pi}{\lambda} \cdot L = n\pi$$

$$\lambda = \frac{2L}{n}$$

↑
integer

and with
 $v = f\lambda$:


$$f = \frac{nv}{2L}$$

Fundamental frequency and harmonics

String of length L
with fixed ends

$$\lambda = \frac{2L}{n}$$

$$f = \frac{nv}{2L}$$



$n=1$: fundamental, first harmonic:

$$\lambda = 2L, f = \frac{v}{2L}$$

$n=2$: 2nd harmonic, 1st overtone:

$$\lambda = L, f = \frac{v}{L}$$

$n=3$: 3rd harmonic, 2nd overtone:

$$\lambda = \frac{2}{3}L, f = \frac{3v}{2L}$$


Examples

A wire has a length of 8m. The speed of waves on the wire is 240m/s. What is the fundamental frequency?

Examples




A particular guitar string has a mass of 3.0 grams and a length of 0.75 m. A standing wave on the string has the shape shown in the figure. The wave has a frequency of 1200 Hz.

- (a) What is the speed of the wave?
- (b) What is the tension of the string?
- (c) The wave on the string produces a sound wave. Does the sound wave have the same frequency, wavelength, or speed as the wave on the string?





Interference

Two or more
traveling waves
superimpose:
interference




Constructive



Destructive

Interference and path length



Two sources, waves emitted in phase. How do waves combine at P?

If crests of both waves arrive at same time:

constructive interference

integer n

$$|L_1 - L_2| = \Delta L = n\lambda$$


If crest of wave 1 arrives at same time as trough of wave 2:

destructive interference

$$|L_1 - L_2| = \Delta L = (n + \frac{1}{2})\lambda$$

Example

Two radio transmitters (**A** and **B**) are 12 m apart. They are driven by the same oscillator (i.e., they emit in phase) and generate waves of wavelength 2 m. How do the waves interfere at point **P** that is 16 m directly in front of source **A**?



Example

Two loudspeakers emit waves of frequency 172 Hz. The speed of sound is 344 m/s. You are 8 m from speaker A. How close can you get to speaker B and have destructive interference?

